

DRUG TESTING PANEL & CUTOFF CONCENTRATIONS

Initial Test Analyte	Initial Test Cutoff Concentration	Confirmatory Test Analyte	Confirmatory Test Cutoff Concentration
Marijuana Metabolites	50 ng/mL	THCA \1\	15 ng/mL
Cocaine Metabolites	150 ng/mL	Benzoylecgonine	100 ng/mL
Opiate Metabolites: Codeine/Morphine \2\	2000 ng/mL	Codeine	2000 ng/mL
		Morphine	2000 ng/mL
6-Acetylmorphine (Heroin)	10 ng/mL	6-Acetylmorphine	10 ng/mL
Phencyclidine	25 ng/mL	Phencyclidine	25 ng/mL
Amphetamines \3\ AMP / MAMP \4\ MDMA \6\ (Ecstasy)	500 ng/mL	Amphetamine	250 ng/mL
		Methamphetamine \5\	250 ng/mL
	500 ng/mL	MDMA	250 ng/mL
		MDA \7\	250 ng/mL
		MDEA \8\	250 ng/mL

\1\ Delta-9-tetrahydrocannabinol-9-carboxylic acid (THCA)

\2\ Morphine is the target analyte for codeine/morphine testing

\3\ Either a single initial test kit or multiple initial test kits may be used, provided the single test kit detects each target analyte independently at the specified cutoff

\4\ Methamphetamine is the target analyte for amphetamine / methamphetamine testing

\5\ To be reported positive for methamphetamine, a specimen must also contain amphetamine at a concentration equal to or greater than 100 ng/mL

\6\ Methylenedioxymethamphetamine (MDMA)

\7\ Methylenedioxyamphetamine (MDA)

\8\ Methylenedioxyethylamphetamine (MDEA)